

[Web](#) [Images](#) [Videos](#) [Maps](#) [News](#) [Shopping](#) [Gmail](#) [more](#)
[abdullah.kawsar@gmail.com](#) | [Scholar Preferences](#) | [My Account](#) | [Sign out](#)


[Advanced Scholar Search](#)
**Scholar**





Results 1 - 20 of about 458. (0.65 sec)

### [The implementation of synchronizing operations in various environments](#)

 H Watson · *Software: Practice and Experience*, 1977 · [intelscience.wiley.com](#)

... If the **interrupt** prevention is located within the scope of the locking variable then the process ... is apparently the fact that two thresholds are erected in front of the **critical region** which leads to ... If it is not available the **threshold** can be simulated by an arrangement shown in Figure 3 ...

[Cited by 10](#) · [Related articles](#) · [All 4 versions](#)

### [\[PDF\] Crossover from superconductivity to a negative magnetoresistance in a Cd-Sb alloy near the localization threshold as the temperature is lowered](#)

[\[PDF\] from issr.ac.ru](#)

 VF Ginzburg, VN Zverev, VM Teslenko · ... *CIC OF PISMA V* ... 1992 · [issr.ac.ru](#)

... It turns out that in the **critical region** the onset temperature of the S transition,  $T_n$ , is ... a correlation between the superconducting response and the conductivity type on the / side of the **threshold** ... By carrying out the heating in increments, one can **interrupt** this process and obtain, in ...

[Cited by 2](#) · [Related articles](#) · [Ed. Direct](#) · [All 4 versions](#)

### [Analysis of phylogenetically reconstructed mutational spectra in human mitochondrial DNA control region](#)

 BA Mityaichuk, IB Ryzikov, VB Barikov · ... *Human genetics*, 2002 · [Springer](#)

... procedure was repeated 10,000 times, a critical value  $P^*$  that separates the right **critical region** of the ... sites with 3 or more mutations which was used as the hotspot **threshold** ( $Sh=3$ ) ... Interestingly, the additional mutations (at positions 16186 and 16192) that **interrupt** the poly-C tract ...

[Cited by 46](#) · [Related articles](#) · [Ed. Direct](#) · [All 4 versions](#)

### [Genes and translocations involved in POF](#)

 D Schlösser, L Herrera, L Cásper · ... *American journal of* ... 2002 · [Wiley Online Library](#)

... follicles after menarche and is progressively depleted until it falls below a **threshold** value and ... If ovarian genes are clustered in the **critical region**, every breakpoint in the region should affect ... The notion that translocations **interrupt** genes to cause POF is further weakened by the ...

[Cited by 53](#) · [Related articles](#) · [Ed. Direct](#) · [All 4 versions](#)

### [Subacute electrical stimulation of the hippocampus blocks intractable temporal lobe seizures and paroxysmal EEG activities](#)

 M Velasco, F Velasco, AL Velasco, S Bolaaga · ... *Epilepsia* 2000 · [Wiley Online Library](#)

... stimulation with rectangular pulses **interrupted** only 1 h/day while EEG recordings were taken. Individual pulses consisted of biphasic, Lilly wave pulses 130 per second in fre- quency, 450 ps in duration, 200-400 pA in amplitude. Amplitude was adjusted for **threshold** or slightly ...

[Cited by 13](#) · [Related articles](#) · [Ed. Direct](#) · [All 4 versions](#)

### [A scheduling scheme for network saturated NT multiprocessors](#)

 J Hansen · ... *Proceedings of the USENIX Windows NT Workshop*, 1997 · [usenix.org](#)

... It is possible to specify that a certain amount of the workload must be performed in a **critical region** ... sharing data protected by locks, the effects of the user thread starvation caused by **interrupt**-routines may ... The fixed **threshold** used to shift to polling is a problem in the case where ...

[Cited by 10](#) · [Related articles](#) · [All 12 versions](#)
[\[PDF\] from usenix.org](#)

### [Development of the concurrent process oriented language 'COL'](#)

 M Mitushin, I Shimoyama · ... *Advanced Robotics*, 1987 · [ingentaconnect.com](#)

... control, (c) priority control, (d) shared variable management, (e) real-time operation, (f) I/O handling, (g) **interrupt** handling, and (h) process state control. ... The **critical region** for mutual exclusion is described using "region" as follows: ...

[Related articles](#) · [All 3 versions](#)

### [Color-induced transition to a nonconventional diffusion regime](#)

 GP Tsironis · ... *Physical review letters*, 1998 · [APS](#)

... by construction, the initial population at  $x=0$ ,  $c(x,0)$ , vanishes throughout the whole **critical region**  $T < T_c$  ... Figure 1 shows that  $\rho(t,0)$  is always semidefinite positive even beyond the **threshold**  $T_c$  where ...  $T/T_c$  on  $\rho(t,0)$ , does not provide the correct slope "and does not **interrupt** this as ...

[Cited by 12](#) · [Related articles](#) · [All 3 versions](#)

### [Electrophysiological and behavioral effects of blockade of the nonspecific thalamo-cortical system](#)

 JE Shansky · ... *Brain Research*, 1987 · [Elsevier](#)

... of varying the temperature of the cryo-probe tip when in the **critical region** of ITP ... are enhanced when any part of the total nonspecific thalamo-cortical system is **interrupted** or blocked ... of the cortex, corresponding to an inhibitory state in that region and a high **threshold** of excitability ...

[Cited by 34](#) · [Related articles](#) · [All 2 versions](#)

### [Role of midbrain central gray in pain-induced defensive boxing of rats](#)

 MA Edwards · ... *Physiology & Behavior*, 1974 · [Elsevier](#)

... lesions which extended in anterior, posterior, or ventral directions from the **critical region** (Fig. ... became slightly less reactive on their second test (mean change in **threshold**: +13.0 ... Similarly,

these lesions interrupted the major descending pathways from the forebrain to the central ...

[Cited by 30](#) • [Related articles](#) • [All 4 versions](#)

### Comparison of phenotype between patients with Prader-Willi syndrome due to deletion 15q and uniparental disomy 15

SE Cassidy, M Forsythe, S Haeger, ... • *Journal of Medical Genetics* ... 1997 • [intscience.wiley.com](#)  
... using highly polymorphic microsatellite markers both within the Prader-Willi Syndrome Critical Region (15q11-q13) ... is that one or both of the breakpoints for the deletion interrupt a gene ... insufficient for clear recommendations, but it would appear that the threshold for clinical ...

[Cited by 118](#) • [Related articles](#) • [All 4 versions](#)

### Percolation in short fibres epoxy resin composites: Conductivity behavior and finite size effects near threshold

F Carmona, P Fruthen, ... • *Solid State Communications*, 1984 • Elsevier  
... conductivity explain the reason being that the experiments have been performed outside the critical region, ... In the vicinity of the percolation threshold where  $\sim 10^{-4}$  to  $10^{-1}$  this leads ... Full, dashed and interrupted lines give hypothetical power law dependences with  $u = 1, 0.88$  ...

[Cited by 45](#) • [Related articles](#) • [All 4 versions](#)

### Catheter ablation of the mitral isthmus for ventricular tachycardia associated with inferior infarction

DJ Wilber, DE Kopp, DO Glascock, N Dennis, ... • *Circulation*, 1995 • *Am Heart Assoc*  
... Conclusions The mitral isthmus contains a critical region of slow conduction in some patients with ... using constant current with a 2-ms pulse duration at twice diastolic threshold ... narrow dimensions, may have been simply the most vulnerable point to interrupt this circumferential ...

[Cited by 111](#) • [Related articles](#) • [All 2 versions](#)

[HTML from ojs.ahajournals.org](#)

### Brain stimulation for neurological and psychiatric disorders, current status and future direction

JY Chang • *Journal of Pharmacology and Experimental Therapeutics* ... 2004 • *ASPE*  
... Anterior and centromedian thalamic stimulations may interrupt epileptiform activity relaying to the cortices thus ... The thalamus is the critical region for relaying neural signals to the cerebral cortex and ... et al., (1982) pointed out that naloxone elevated the stimulation threshold in the ...

[Cited by 16](#) • [Related articles](#) • [All 2 versions](#)

[HTML from aspejournals.org](#)

### Conduction in anisotropic disordered systems: Effective-medium theory

J Flourens • *Physical Review B*, 1974 • *APB*  
... It has been successfully applied to percolation models (outside the threshold region) and ... on networks with  $20 \times 20$  nodes is remarkably good, except in a critical region near  $p$  ... This boundness of the anisotropy  $Q$  has important consequences for the interrupted strand model ...

[Cited by 42](#) • [Related articles](#) • [All 2 versions](#)

### Neurons of rostral ventrolateral medulla mediate somatic pressor reflex

Rt. Stometta, SF Morrison, ... • *American Journal of Physiology*, 1989 • *Am Physiological Soc*  
... that it is mediated by spinoreticular afferents traveling in the contralateral spinal cord, and that the C1 adrenergic area of the RVLM is the critical region for the ... stimulation (stimulus trace) is a 5-s train of square-wave pulses (1.0 ms, 20 Hz) at a stimulus current 5 times threshold (see ...

[Cited by 51](#) • [Related articles](#) • [All 3 versions](#)

### Toward a molecular understanding of congenital heart disease

RM Payne, MC Johnson, JW Grant, ... • *Circulation*, 1995 • *Am Heart Assoc*  
... resultant mutant phenotype will provide clues to the normal function of the interrupted gene ... In the threshold model, which some have proposed to explain human CHD, 2 multiple ... combined with the trisomy 16 (T16) mouse model have narrowed the critical region for endocardial ...

[Cited by 111](#) • [Related articles](#) • [All 3 versions](#)

[HTML from ahajournals.org](#)

### Phase transitions in relational learning

A Gloriano • *Machines Learning*, 2000 • Springer  
... From figures 8(a)-(d) we would predict that formula  $\phi_1$  should be easy to match for all the examples, whereas  $\phi_2$  is likely to require a high computational cost to be matched, because most examples lay in the critical region. ...

[Cited by 56](#) • [Related articles](#) • [All 3 versions](#)

[PDF from unigal.it](#)

### [HTML] Disruption of a novel member of a sodium/hydrogen exchanger family and DOCK3 is associated with an attention deficit hyperactivity disorder-like phenotype

MG Da Silva, K Elliott, RH Doherty, ... • *Journal of Molecular Biology* ... 2003 • [pubs.bmj.com](#)  
... assembled using co-segregation analysis of EST markers along the BAC clones in the critical region and sequence ... The gene interrupted at the p arm, DOCK3, is a member of the GDM family ... Although the lod score did not exceed the QTL threshold for significance, its location within ...


[Cited by 30](#) • [Related articles](#) • [All 2 versions](#)

[HTML from bmc.com](#)

### Approximate analysis of distributed semi-hard real-time systems

D Maniescu • *Automatic Control IEEE Transactions on*, 2002 • [ieeexplore.ieee.org](#)  
... and S. B. Gershwin, "The uncertainty threshold principle: Some fundamental limitations of optimal decision making under ... CS, is a service center corresponding to execution in critical region i ... A process with a higher priority cannot interrupt the execution of a process with lower ...

[Cited by 12](#) • [Related articles](#) • [All 3 versions](#)

 [Create email alert](#)